

COLORING ABELIAN GROUPS

P.D. JOHNSON

Department of Mathematics, American University of Beirut, P.O. Box 11-0236, Beirut, Lebanon

Received 9 January 1981

A sufficient (and necessary, if $n=2$) condition for the existence of a particular kind of n -coloring of an abelian group is given, and applied to show that (a) the real line is colorable with two colors so that the distance 1 is forbidden for one color, and the distance $s > 0$ for the other, or so that both 1 and s are forbidden for both colors, if and only if s is not the ratio of an odd and an even integer; (b) the chromatic number of Q^2 and Q^3 is 2, but that of Q^n is greater than 2 for $n > 3$.

The object of this paper is mainly to give and apply a necessary and sufficient condition for the two-colorability of certain graphs associated with abelian groups. This condition is, of course, equivalent to the condition of non-existence of odd cycles in the graph, but has an algebraic character that makes it easier to apply in certain situations.

G will be an abelian group. An n -coloring of G will be a partition of G into n sets.

Definition. Suppose $h \in G$; $h \neq 0$. An n -coloring of G is h -alternating if and only if, for each $g \in G$, the group elements $g + mh$, $0 \leq m \leq n-1$, are all colored differently.

A subset $B \subseteq G$ is h -forbidden if and only if, for each $g \in G$, $g \in B$ implies $g + h \notin B$. A coloring of G is h -forbidden if and only if each of the sets of the partition is h -forbidden.

If $S \subseteq G$, a coloring of G is S -alternating, or S -forbidden, if and only if it is h -alternating, respectively h -forbidden, for each $h \in S$.

A subset $S \subseteq G$ is weakly n -free if and only if $\sum_{h \in S} m_h h = 0$ implies $\sum_{h \in S} m_h \equiv 0 \pmod{n}$. [Here, and later, at most finitely many of the integers m_h are non-zero. The word 'weakly' is used because the sum of the m_h is required to be divisible by n , not each m_h separately.]

It will be useful to note that if a coloring is h -alternating, then it is h -forbidden; if a set $B \subseteq G$ is h -forbidden, then it is $-h$ -forbidden; if an n -coloring is h -alternating, then $g + mh$ and $g + kh$ have the same color if and only if $m \equiv k \pmod{n}$. In the case $n = 2$, this last observation may be extended: if $S \subseteq G$, then

for any S -alternating two-coloring, g and $g + \sum_{h \in S} m_h h$ have the same color if and only if $\sum_{h \in S} m_h$ is even.

The case $n = 2$ is also distinguished by the following.

Proposition. *A two-coloring B, R of G is $\{h, z\}$ -alternating if and only if B is h -forbidden and R is z -forbidden.*

Proof. The 'only if' assertion is trivial. Suppose B is h -forbidden and R is z -forbidden. If $g, g+h \in R$, then $g+z, g+z+h \in B$; but this contradicts the assumption that B is h -forbidden. Consequently, at least one of $g, g+h$ is in B , and thus exactly one, since B is h - (and therefore $-h$ -) forbidden.

Thus B, R is h -alternating, and, likewise, z -alternating.

[*Remark.* This proposition is a slight improvement over the easy observation that a two-coloring is S -forbidden if and only if it is S -alternating.]

Theorem. *Suppose $S \subseteq G$, and $n > 1$. If S is weakly n -free, then there is an S -alternating n -coloring of G . If $n = 2$, or if S is a singleton, the converse holds; the existence of an S -alternating n -coloring implies that S is weakly n -free.*

Proof. Suppose S is weakly n -free. Let H denote the subgroup generated by S . Color H by letting

$$B_k = \left\{ \sum_{h \in S} m_h h \mid \sum_{h \in S} m_h \equiv k \pmod{n} \right\}, \quad k = 0, \dots, n-1.$$

The B_k are mutually disjoint by the assumption that S is weakly n -free. Extend the coloring to all of G by coloring the co-sets of H similarly.

The proof of the converse in the special cases mentioned is easy, by previous remarks.

Corollary 1. *Suppose S_1 and S_2 are non-empty subsets of G . There is a two coloring B, R of G such that B is S_1 -forbidden and R is S_2 -forbidden if and only if $S_1 \cup S_2$ is weakly two-free.*

Proof. This is an easy corollary of the Proposition and the Theorem.

Corollary 2. *Suppose $s > 0$. There is a two-coloring of the real line (with red and blue, say), such that no two red points are 1 apart, and no two blue points are s apart, if and only if $s \notin K = \{p/q \mid p, q \text{ are positive integers and } p+q \text{ is odd}\}$.*

Proof. The conclusion follows from Corollary 1 and the observation that $\{1, s\}$ is weakly 2-free in the additive group of reals if and only if $s \notin K$.

The chromatic number of a subset of a Euclidean space is the smallest number

of colors needed to color the set so that no two points a distance 1 apart are the same color (the distance 1 is forbidden).

Corollary 3. *The chromatic number of Q^2 and Q^3 is 2, and of Q^4 is greater than 2.*

Proof. To show that the chromatic number of Q^2 is 2, it suffices to show that $S = \{(x, y) \in Q^2 \mid x^2 + y^2 = 1 \text{ and } x = 1 \text{ or } y > 0\}$ is weakly 2-free (in the additive group). This is essentially a consequence of the fact that if (a, b, c) is a primitive Pythagorean triple, then one of a, b is odd, the other even, and c is odd. Thus, if $\{(1, 0), (0, 1), (a_1/c_1, b_1/c_1), \dots, (a_k/c_k, b_k/c_k)\}$ is a finite subset of S (into which it doesn't hurt to throw $(1, 0)$ and $(0, 1)$), with (a_i, b_i, c_i) being relatively prime Pythagorean triples, and

$$n(1, 0) + r(0, 1) + \sum_i m_i(a_i/c_i, b_i/c_i) = (0, 0),$$

then, multiplying by c , the product of the c_i , an odd number, we have

$$n + \sum m_i a_i \equiv 0 \pmod{2} \quad \text{and} \quad r + \sum m_i b_i \equiv 0 \pmod{2},$$

so

$$n + r + \sum m_i(a_i + b_i) \equiv n + r + \sum m_i \equiv 0 \pmod{2},$$

which was to be shown.

The proof for Q^3 is similar. The point there is that in any primitive Pythagorean quadruple (a, b, c, d) , d and exactly one of a, b, c must be odd.

In Q^4 , we have $3 \cdot [\frac{1}{6}(1, 1, 3, 5)] - 1 \cdot [\frac{1}{2}(1, 1, 1, 1)] - (0, 0, 1, 0) - 2(0, 0, 0, 1) = (0, 0, 0, 0)$, while $3 - 1 - 1 - 2$ is odd, which shows that the unit sphere in Q^4 is not weakly 2-free, which means that Q^4 cannot be colored with two colors so that the distance 1 is forbidden.

Remarks. (i) In Corollary 2, if $s = p/q$, p, q odd positive integers, then the coloring called for could be obtained by alternating colors on the intervals

$$\left[\frac{n}{q}, \frac{n+1}{q} \right), \quad n = 0, \pm 1, \dots$$

However, if s is irrational, then for every 2-coloring forbidding 1 and s , both colors are dense on the line. To see this, note that $A = \{m + ns \mid m, n \text{ integers}\}$ is dense on the line, and the sets $2A, 2A + 1$ are each monochromatic, of different colors.

(ii) Again regarding Corollary 2, the theorem really says that the line can be 2-colored forbidding all distances in a set $S \subseteq (0, \infty)$ if and only if S is weakly 2-free. The existence of weakly 2-free sets and Zorn's Lemma imply the existence of maximal weakly 2-free sets; it is easy to see that any such set must span the

reals, considered as a vector space over Q , and thus must have cardinality 2^{\aleph_0} . It is generally well known that 2^{\aleph_0} distances are forbiddable in a 2-coloring of the line; the power of the theorem in this context is not in the rederivation of this result, but in that the theorem describes which sets of distances are forbiddable.

As a possible application, note that if a weakly 2-free set $S \subseteq (0, \infty)$ could be found with the property that $x^2 + y^2 = 1$, $x, y \geq 0$ implies $x \in S$ or $y \in S$, then we could 4-color the plane so that the distance 1 is forbidden for all colors; just color the line with r and b so that all distances in S are forbidden, and then color the plane with the four colors (r, r) , (r, b) , (b, r) , (b, b) in the obvious way. Thus the existence of such a set S would imply that the chromatic number of the plane is 4 (see [2]), although its non-existence would not imply the contrary. However, the author does not know whether or not such an S exists.

(iii) The result that the chromatic number of Q^2 is 2 is known, and a brief proof is given in [2, Theorem 1].

(iv) Consider the following problem scheme: you are given a subset A of a Euclidean space, and k sets of distances S_1, \dots, S_k , and are asked to color A with k colors c_1, \dots, c_k so that all distances in S_i are forbidden for the color c_i , $i = 1, \dots, k$. The special cases in which the S_i are singletons are of particular interest. The general problem scheme can be formulated in the terminology of [1], but doesn't seem to be a Euclidean Ramsey problem scheme in any obvious way.

Woodall gives a proof of a theorem of Raiskii [2, Theorems 2 and 3] which certainly implies that if A is Euclidean n -space, and $k = n + 1$, $n > 1$, then the problem has no solution for any non-empty S_1, \dots, S_k . On the other hand, D. Greenwell, in unpublished work, has shown that if $k = 3$ and A is the line, then the problem can be solved for every sequence S_1, S_2, S_3 of singletons (and it is a consequence of Corollary 2 that 3 is the smallest k of which this can be said). The Proposition of this paper implies that if A is an additive subgroup of a Euclidean space, then, for $k = 2$, the problem can be solved for S_1, S_2 if and only if a 2-coloring can be found for which all the distances in $S_1 \cup S_2$ are forbidden for both colors; the theorem implies that such a coloring can be found if and only if $\{x \in A \mid |x| \in S_1 \cup S_2\}$ is weakly 2-free (here $|\cdot|$ stands for the Euclidean norm).

Specific unsolved problems of the type described swarm by the gazillions, but at least two seem to be indicated by the results cited:

(a) In the case when A is the line and $k = 3$, extend Greenwell's result away from singletons, and the consequences of this paper away from $k = 2$, by finding simple conditions on sets S_1, S_2, S_3 of distances for the solvability of the problem.

(b) When A is the plane, find the smallest k , if any exists, such that the problem can be solved for any sequence S_1, \dots, S_k of singletons.

Acknowledgements

Thanks, but no blame, are due to Don Greenwell, Dean Hoffman, and David Makinson for conversations from which the author benefited, on the subject of this paper.

References

- [1] Erdős, Graham, Montgomery, Rothschild, Spencer and Straus, Euclidean Ramsey Theorems I, J. Combin. Theory (A) 14 (1973) 341–363.
- [2] D.R. Woodall, Distances realized by sets covering the plane, J. Combin. Theory (A) 14 (1973) 187–200.